Guidelines for use of personal protective equipment in relation to Covid-19:

Developed by expert Team of NMC and Government of Nepal with reference from WHO

- A. For Aerosol Generating procedures: Dental procedures, bronchoscopy, Upper GI Endoscopy, ENT procedures, Nebulization, Intubation of a patient, CPR, Non-invasive ventilation, endotracheal suctioning, when obtaining nasopharyngeal or oropharyngeal swab, etc. in Covid-19 suspected or confirmed cases health personnel need to use the following protective equipment: Category I PPE:
 - a. N-95 mask
 - b. Goggles or visor
 - c. Gloves (loose gloves acceptable)
 - d. Water resistant OR standard disposable gowns
 - e. Cap: Regular disposable
- B. For Non aerosol generating covid-19 suspected or confirmed patients: Health personnel need to use the following protective equipment: Category II PPE:
 - a. Surgical mask (seal the top edge with tape)
 - b. Goggles or visor
 - c. Gloves (loose gloves acceptable)
 - d. Water resistant or standard disposable gowns
 - e. Cap: Regular disposable
- For Physicians examining the patients in the fever/screening clinics the following PPE is recommended: Category II PPE
 - a. Surgical mask, (seal the top edge with a tape)
 - b. Goggles or visor
 - c. Water resistant or standard disposable gowns
 - d. Regular disposable Cap
 - e. Gloves (loose gloves acceptable)

Note: For staff involved in preliminary screening not involving direct contact, maintain spatial distance of at least 1meter, no PPE required; Use surgical mask if maintaining distance not always possible.

- D. For escorts or drivers, the following PPE is recommended: Category III PPE:
 - a. Surgical masks
 - b. Gloves
 - Gowns (if physical contact expected)
 Follow hand hygiene and keep minimum 3 to 6 feet distance from the patient
- E. For Laboratory staff: category II or III PPE; depending upon the chance of splash:
 - d. surgical mask
 - e. Gown
 - f. Loose Gloves
 - g. Eye protection (if risk of splash)
- F. For all staff, including health care workers involved in any activity that does not involve contact with COVID-19 patients and working in other areas of patient transit (e.g. wards, corridors). No PPE required.

For Everyone:

- o Maintain 3-6 feet distance while visiting patients, if no need to touch the patient.
- Mandatory hand-hygiene after each use of PPE and between patients.
- Mandatory surface cleaning of bed or furniture with 0.5% Chlorine disinfectant (Virex*
 or similar) between each patient in OPD or in an inpatient setting.





Cololà